



## **The haematophagous arthropods (Animalia: Arthropoda) of the Cape Verde Islands: a review**

**Elves Heleno Duarte<sup>1</sup>**

Keywords: Arthropods, arthropod-borne diseases, bloodsucking, hematophagy, Cape Verde Islands

### ABSTRACT

Arthropoda is the most diverse phylum of the animal kingdom. The majority of bloodsucking arthropods of public health concern are found in two classes, Arachnida and Insecta. Mosquitoes, ticks, cattle flies, horseflies and biting midges are the main hematophagous groups occurring in the Cape Verde Islands and whose role in infectious disease transmission has been established. In this literature review, the main morphological and biological characters and their role in the cycle of disease transmission are summarized.

### RESUMO

Os artrópodes constituem o mais diverso entre todos os filões do reino animal. É na classe Arachnida e na classe Insecta que encontramos a maioria dos artrópodes com importância na saúde pública. Os mosquitos, os carrapatos, as moscas do gado, os tabanídeos e os mosquitos pólvora são os principais grupos hematófagos que ocorrem em Cabo Verde e possuem clara associação com a transmissão de agentes infecciosos. Nesta revisão da literatura apresentamos os principais caracteres morfológicos e biológicos e o seu papel no ciclo de transmissão de doenças.

---

<sup>1</sup> Direcção Nacional da Saúde, Ministério da Saúde, Avenida Cidade de Lisboa, C.P. 47, Praia, Republic of Cape Verde; [ehelgam@gmail.com](mailto:ehelgam@gmail.com)

## INTRODUCTION

With over a million described species, Arthropoda is the most diverse and species rich clade of the animal kingdom. Five main taxonomic groups are usually recognized: the extinct Trilobitomorpha and the extant Chelicerata, Miriapoda, Hexapoda and Crustacea (Ruppert & Barnes 1994). Because they represent an important source of food, transmit numerous infectious agents and include an array of agricultural pest species, arthropods are widely studied (Chown & Nicolson 2004). Haematophagous arthropods occur in two major groups, i.e. Arachnida (Chelicerata) and Insecta (Hexapoda) (Fig. 1), and are vectors of pathogens worldwide. So far, more than 16,000 haematophagous species have been identified, of

which *ca.* 500 are strongly associated with the transmission of infectious agents (Grimaldi & Engel 2005, Lehane 2005). It has been estimated that of infectious diseases worldwide, about 17% are vector-borne. Unfortunately, vaccines for most of these diseases are not available. Therefore, increased emphasis on vector control strategies is required, based upon the selection of proven intervention methods tailored to biological characters and ecological circumstances of local vectors (WHO 2004).

The terrestrial arthropod fauna of the Cape Verde Islands was reviewed by van Harten (1993), while a summary update was recently provided by Arechavaleta *et al.* (2005). In Cape Verde, several vector-borne diseases occur.




Fig. 1. Main groups of haematophagous arthropods occurring in the Cape Verde Islands. Modified after Lehane (2005) and Estrada-Peña *et al.* (2010).

Many of these are emerging and/or re-emerging as a result of ecological and environmental changes that favour increased vector densities (Gratz 1999). In order to reduce the chance of (re)emergence of arthropod-transmitted diseases, knowledge of local vector populations is crucial in tracking any changes in their biology. This review was conducted in order to draw attention

to the importance of bloodsucking arthropods in the Cape Verde archipelago and to better understand their role in the most common infectious agent transmissions. The result will hopefully be useful in making policy decisions and in formulating new strategies in the fight against vector-borne diseases in the Cape Verde Islands.

## METHODS

The literature was scanned using the PubMed, Google Scholar and SCIELO databases from January 2013 to December 2013. Combinations of the following key words were used: arthropods, arthropod-borne disease, blood-sucking, haematophagy and Cape Verde Islands. Three different languages (English, French, Portuguese) were used to obtain results. Unpublished reports in national institutions (e.g. Ministry of Health, Ministry of Rural Development, National Library, National Historical

Archive) were also scrutinized. Relevant references were organized in a spreadsheet, but only those references actually mentioning bloodsucking arthropods were maintained. Using the same databases, the medical and/or veterinary importance of the references was assessed. Medical and/or veterinary importance was allotted when at least one infectious agent was isolated in the wild or when experimental infection was successfully realized in the laboratory.

## RESULTS

The database search resulted in 22 publications (articles, books, reports) mentioning the presence of haematophagous arthropods in the Cape Verde Islands. Of extant bloodsucking arthropods, only Lepidoptera (represented by a single haematophagous species in Southeast Asia) do not occur in the archipelago. Only one study (on mosquitoes) was conducted exclusively by national researchers (Duarte *et al.* 2012). Also focusing on mosquitoes, six publications were published by a combined team of national and foreign workers (Appendix 1). The remaining studies were conducted solely by foreign workers and dealt with ticks, cattle flies, horseflies, biting midges, blackflies, mites, fleas and lice. Publications appeared in 19 journals and other sources (Appendix 1). Up until now, a role in the disease transmission cycle has been confirmed for three taxonomic groups in the Cape Verde archipelago (see group descriptions below).

### TICKS (ARACHNIDA: ACARI: IXODIDA)

Ticks are mandatory ectoparasites that parasitize a variety of vertebrates and cause direct and indirect financial loss (Parola & Raoult 2001, Estrada-Peña *et al.* 2010). This large group is subdivided into three families: i) Ixodidae (hard

ticks), with over 700 species, including the most important vectors; ii) Argasidae (soft ticks), comprising *ca.* 200 species; iii) Nuttalliellidae, with a single species (*Nuttalliella namaqua*), exclusively found in southern Africa (Parola & Raoult 2001, Basu *et al.* 2012).

Being arachnids, ticks can be easily distinguished from insects by having three pairs of appendices during the immature stage and four pairs as adults, by having the mouthpart transformed into chelicerae and by the absence of wings (Randolph 1998). Only few studies on ticks have been conducted in the Cape Verde Islands (Tendeiro 1954, Meira *et al.* 1957, Kirchner *et al.* 2008, Götsch *et al.* 2009, Gómez-Díaz *et al.* 2012). All except the study by Gómez-Díaz *et al.* (2012) were conducted fully or partially in Santiago Island. Recent studies showed domestic animals to be highly parasitized and it was recommended to prevent transportation of animals (especially dogs) from Cape Verde to Europe (Kirchner *et al.* 2008, Götsch *et al.* 2009). *Ripicephalus sanguineus* was the only species found in recent studies, although other species such as *Amblyomma variegatum*, *Margaropus decoloratus* and *Hyalomma* sp. had previously been reported (Tendeiro 1954, Meira *et al.* 1957), all of them

Ixodidae. The most recent study of ticks conducted in the archipelago (Gómez-Díaz *et al.* 2012) dealt with diversity and genetic structure of *Ornithodoros capensis*, a parasite of seabirds.

#### LIFE CYCLE

Ticks have a complex life cycle, which – depending on family, species and environmental parameters – may take 2-3 years (Parola & Raoult 2001). Their feeding behaviour is also complex. Hard ticks need a long time to feed (2-15 days) and feeding only takes place once during each stage (larva, nymph and adult). Soft ticks consume several meals per stage but, unlike hard ticks, these may take from a few minutes to a few hours (Vial 2009). Approximately 75% of a hard tick's life cycle is spent while being attached to an animal (Parola & Raoult 2001), whereas soft ticks only attach to animals to feed (Vial 2009).

#### MEDICAL AND VETERINARY IMPORTANCE

African swine fever (ASF) is a highly contagious and extremely deadly disease in domestic pigs (FAO 2000) and is one of the most important tick-borne ailments in Cape Verde. Ticks of the genus *Ornithodoros* play an important role as vector of the disease (Basto *et al.* 2006). Although there is no consensual view on their precise role in ASF transmission, the persistence of the disease after several veterinary interventions and absence of transmission during various months (Penrith 1998) suggests that vectors or pigs (or both) act as reservoirs.

No human diseases transmitted by ticks are known to occur in the archipelago. Lyme disease is an illness caused by the spirochete *Borrelia burgdorferi s.l.* and is transmitted by hard ticks of the genus *Ixodes* (Karami 2012), but has not been found in Cape Verde so far. Beyond their implication in disease transmission, ticks themselves pose problems to both man and animals because they cause various harmful side-effects to the host, ranging from anaemia caused by massive infestation to allergy due to the inoculation of saliva during blood meals (Lehane 2005).

#### MOSQUITOES (INSECTA: NEMATOCERA: CULICIDAE)

Mosquitoes are dipteran insects belonging to the family Culicidae, in which three subfamilies are recognized: Anophelinae, Culicinae and Toxirhichitinae (Consoli & Lourenço de Oliveira 1994). Diptera have only one pair of wings, the

forewings, while the hindwings are reduced to dumbbell-shaped knobs called halteres. They have long legs and antennae, chipper-shaped mouthparts adapted to suction and generally show marked sexual dimorphism (Consoli & Lourenço de Oliveira 1994, Harbach 2007). In the Cape Verde archipelago, 11 species of mosquitoes occur, representing two subfamilies, i.e. Anophelinae (two species in one genus) and Culicinae (nine species in three genera) (Ribeiro *et al.* 1980, Alves *et al.* 2010, in press). Of these, about half is involved in the transmission of infectious agents, particularly *Anopheles arabiensis*, *Aedes aegypti* and two members of the *Culex pipiens* complex (*C. p. pipiens* and *C. p. quinquefasciatus*) (Alves *et al.* 2010).

#### LIFE CYCLE

During their life cycle, mosquitoes pass through four stages: eggs, larvae, pupae and adults, of which the first three are aquatic (Consoli & Lourenço de Oliveira 1994, Lehane 2005). The larvae feed mostly on organic particles in water, while pupae only use the energy stored during the larval stage. Adult mosquitoes are terrestrial, this being the stage of reproduction and dispersion. Males feed exclusively on plant fluids, while females need animal (including human) blood for the maturation of their eggs (Consoli & Lourenço de Oliveira 1994).

#### MEDICAL AND VETERINARY IMPORTANCE

In Cape Verde, mosquitoes have been identified as vector of several infectious agents that cause malaria, yellow fever, lymphatic filariasis and, more recently, dengue (Franco & Menezes 1955, Ribeiro *et al.* 1980; Alves 2004, WHO 2009).

*A. arabiensis* is the only member of the *A. gambiae* complex occurring in Cape Verde (Cambournac *et al.* 1982, Diallo 2003, Alves *et al.* 2010, Dia *et al.* 2011). In addition to being the only vector of malaria, it was also the vector of *Wulchereria bancrofti*, the infectious agent causing lymphatic filariasis (Franco & Menezes 1955). Since the 1950s, no new cases of lymphatic filariasis have become known in Cape Verde and the disease has seemingly been eradicated in the islands. In a recent study conducted in all inhabited islands, no cases were diagnosed (Benzerroug 2005). Before the 1950s, the annual incidence of malaria was more than 100 cases/1000 inhabitants (Rodriguez *et al.* 2012), but currently only limited and localized transmission occurs in two (Santiago and Boa Vista) of the 10 islands (WHO 2012).

Epidemic dengue fever occurred in Cape Verde in 2009 when *ca.* 21,000 cases were reported (WHO 2009, Monteiro 2010), mainly in Santiago and Fogo Islands. *A. aegypti*, the only vector of dengue described in the archipelago (Alves *et al.* 2010), was also the only vector of yellow fever, being resistant to DDT 4% and also suspected to be resistant to propoxur (Dia *et al.* 2012). During the dry season, it takes advantage of household water containers for its reproduction, thus maintaining high densities over the reproductive period (Duarte *et al.* 2012, 2013). Using experimental infection techniques, it has been shown that *A. aegypti* (ssp. *formosus*) from Santiago Island has a moderate ability to transmit dengue virus serotype 3, but a high susceptibility of becoming infected with and to transmit chikungunya (CHIKV) and yellow fever virus (Vazeille *et al.* 2013).

Despite their role in the transmission of several infectious agents in other countries, the two members of the *C. pipiens* complex that occur in Cape Verde have as yet not been associated with infectious agent transmission. Elsewhere, these taxa are instrumental in the transmission of West Nile virus, *Wulchereria bancrofti*, Rift Valley fever viruses, encephalitis viruses and others (Turell 2012). In Cape Verde, *C. p. quinquefasciatus* was first documented in 1950 (although the presence of *C. pipiens s.l.* had already been reported in 1947), while the occurrence of *C. p. pipiens* was established in 1977 (Ribeiro *et al.* 1980). Based on morphological studies of the male genitalia, Ribeiro *et al.* (1980) identified hybrids *C. p. pipiens* x *C. p. quinquefasciatus*. This was subsequently confirmed by molecular analysis (Alves *et al.* 2010, Gomes *et al.* 2012). These hybrids have been shown to have the ability to enhance arbovirus transmission in areas where they occur (Gomes *et al.* 2012).

#### **CATTLE FLIES (INSECTA: BRACHYCERA: MUSCIDAE)**

Cattle flies are bloodsucking ectoparasites of mammals (especially cattle) in the genus *Stomoxys*. They are similar to houseflies *Musca domestica*, but the distinguishing character is the cattle flies' mouthparts, which are adapted to bloodsucking (Zumpt 1973). Both males and females feed on blood. Three species occur in Cape Verde, i.e. *Stomoxys calcitrans*, *S. niger* and *S. sitiens* (Arechavaleta *et al.* 2005), of which only *S. calcitrans* has anthropophilic preferences.

#### **LIFE CYCLE**

During their life cycle, cattle flies go through four stages: egg, larvae, pupae and adult (Lehane 2005). The eggs are laid in groups of 40 to 80. Hatching occurs approximately 24 hours after laying, while larval development time depends on temperature and other environmental conditions. After the last instar, larvae move to dry areas for pupation. Adults live for about 30 days, with males on average living slightly longer than females (Zumpt 1973, Lehane 2005).

#### **MEDICAL AND VETERINARY IMPORTANCE**

Cattle flies are characterized by having interrupted blood meals and they can bite several hosts during the course of the same feeding round. This has important epidemiological consequences (Zumpt 1973). Therefore, their economic damage is categorized as either direct or indirect. Direct damage is inflicted by blood spoliation, decrease in immune defense (inducing latent diseases), production loss, diminished weight, etc. Indirect damage is caused by the transmission of viruses, bacteria and other infectious agents (Zumpt 1973, Lehane 2005).

*S. calcitrans* is a pest species with a worldwide distribution, known for disturbing cattle and causing considerable losses (Lehane 2005). The species can also transmit trypanosomes, mainly *Trypanosoma equinum* in Neotropical countries and *T. evansi* (which causes severe disease in horses and dogs and less severe illness in cattle) and it has a secondary role in the transmission of the infectious agent causing African trypanosomiasis or sleeping sickness (Lehane 2005). The only reported link between *S. calcitrans* and disease in Cape Verde occurred in the past, when its larvae caused myiasis among humans (Azevedo & Moreira 1946).

#### **HORSEFLIES (INSECTA: BRACHYCERA: TABANIDAE)**

Horseflies are robust insects (adults: 5-25 mm) with a cosmopolitan distribution. The males feed on plants, while the hematophagous females also feed on nectar (Middlekauff & Lane 1980, Lehane 2005). Their head is larger than the thorax, the mouthparts are of the chipper/sucking type and they have long antennae. The Tabanidae comprise more than 4,300 described species in more than 130 genera and three subfamilies (Tabaninae, Chrysopsinae, Pangoninae) of which the first two are the epidemiologically more important (Lehane 2005). *Atylotus agrestis* is



probably the only species that occurs in the Cape Verde Islands (Arechavaleta *et al.* 2005).

#### LIFE CYCLE

The life cycle of horseflies includes eggs, larvae (with 6-13 stages), pupae and adults. Egg laying occurs in aquatic environments and eggs hatch 2-3 days after laying. Larvae also need humid environments to survive; they are carnivorous and feed on small invertebrates. Horseflies can remain at the larval stage for up to two years before transforming into pupae. After 1-3 weeks, adults emerge and live for about two months. Mating occurs soon after emergence and females lay their eggs only after having consumed blood meals (Middlekauff & Lane 1980, Lehane 2005).

#### MEDICAL AND VETERINARY IMPORTANCE

Horseflies possess some characteristics that favour the transmission of infectious agents: only few species are autogenous (most require a blood meal for egg maturation), they are telmophagous (skin deceleration during blood meal), they require a fair amount of blood (and thus have a long engorgement time) and they interrupt their meal due to being chased off because of their painful bite, thus seeking another host (Middlekauff & Lane 1980, Lehane 2005). They may transmit a large variety of infectious agents, including bacteria, viruses, protozoa, filariae and others. Anthrax, anaplasmoses, Q fever, trypanosomiasis, filariasis, encephalitis and African swine fever are some of the diseases transmitted (Lehane 2005). Because some of these diseases occur in Cape Verde and due to the fact that *Atylotus agrestis* is associated with the transmission of some infectious agents (Desquesnes & Dia 2003a, 2003b), further studies are needed to clarify the role of this species as a vector in the archipelago.

#### BITING MIDGES (INSECTA: NEMATOCERA: CERATOPOGONIDAE)

Biting midges are small (1-4 mm) flies of the family Ceratopogonidae, having compound eyes, chipper-shaped mouthparts, short legs and the abdomen divided into 10 segments (Mellor *et al.* 2000). With the exception of New Zealand, Patagonia, the Hawaiian Islands and the polar regions, they have a worldwide distribution. The genus *Culicoides* includes ca. 1,400 species of which 96% engage in bloodsucking (females only). They parasitize mammals (including humans) and birds (Mellor *et al.* 2000, Zimmer *et al.* 2008). *C. imicola* – the main vector of

African horse sickness virus (AHSV) and Bluetongue virus (BTV) in Africa – *C. schultzei* and *C. nivosus* occur in the Cape Verde archipelago (Boorman & van Harten 1992).

#### LIFE CYCLE

The *Culicoides* life cycle includes eggs, four larval stages, pupae and adults. The immature stages require humid places to survive (Kettle 1977, Mellor *et al.* 2000). Breeding sites are similar to those of mosquitoes. Eggs are laid at the substrate surface and, depending on species and environmental conditions, hatching occurs 2-7 days after laying (Mellor *et al.* 2000). Larvae feed on vegetal debris, but some species are predators. Pupae can be found moving free in the water or fixed on the substrate. Depending on the species, adults are active during daylight or twilight, possess only limited capacity for flight and dispersal and are generally passive (Kettle 1962, 1977, Mellor *et al.* 2000).

#### MEDICAL AND VETERINARY IMPORTANCE

Worldwide, more than 50 arboviruses have been isolated from *Culicoides*, sometimes playing a secondary role in the transmission cycle (Mellor *et al.* 2000). Many species transmit infectious agents causing diseases in animals, but only few of them in humans. Among infectious agents transmitted, Rift Valley fever (RVF) virus, African horse sickness (AHS) virus, bluetongue virus (BTV), equine encephalitis viruses and epizootic hemorrhagic disease (EHD) virus are some examples (Mellor *et al.* 2000, MacLachlan & Guthrie 2010). Two of these agents, AHSV and BTV (Orivirus, Reoviridae), cause diseases of significant international impact and have been reported in the Cape Verde Islands (Sellers *et al.* 1977, Boorman & van Harten 1992). AHSV is a non-contagious disease that causes 90% mortality in infected horses and has been introduced in Cape Verde from Senegal (Sellers *et al.* 1977, MacLachlan & Guthrie 2010). Nine serotypes of AHSV that occur in Africa are transmitted by *C. imicola* and *C. bolitinos* (MacLachlan & Guthrie 2010).

#### OTHER TAXA

A single species of black fly (Nematocera: Simuliidae), *Simulium ruficorne*, occurs in the Cape Verde archipelago (Arechavaleta *et al.* 2005). Worldwide, there are about 1,800 species of black flies in 25 genera, of which four are of public health concern: *Austrosimulium*, *Cnephia*, *Prosimulium* and *Simulium* (Lehane 2005).

These black flies transmit *Onchocerca volvulus*, which causes onchocerciasis in Africa, but in Cape Verde *S. ruficornis* has as yet not been shown to be a vector of infectious agents.

Not surprisingly, *Cimex hemipterus* (Heteroptera: Cimicidae), the common bedbug, also occurs in Cape Verde (van Harten 1993, Arechavaleta *et al.* 2005). Both sexes are hematophagous and preferably bite at night (Lehane 2005). Although they are suspected of transmitting infectious agents, the role of bedbugs in spreading them is not clear and there is no clear evidence for their involvement in the transmission of disease agents (Delaunay *et al.* 2011).

Both Hippoboscidae (louse flies) and Oestridae (botflies) occur in the Cape Verde archipelago (van Harten 1993, Arechavaleta *et al.* 2005). Van Harten (1993) cited three species for the archipelago: *Hippobosca rufipes*, *H. equine* and *Olfersia aenescens*. Worldwide, more than 200 species have been described and several of them have been implicated in the transmission of infectious agents (e.g. Rahola *et al.* 2011). *Oestrus ovis* is the only species of botfly occurring in Cape Verde and it has been implicated in causing myiasis in several species elsewhere in the world (Denion *et al.* 2004).

Although many species of mites (Siphonaptera) have been confirmed to occur in the Cape Verde Islands (Mahunka 1991, Arechavaleta *et al.* 2005, Haitlinger 2009), only Erythraeidae (*Leptus salicus*, *L. korneli*,

*Erythraeus capeverdensis*) may have some degree of hematophagous habits. Although few data are available for these species, it has been shown elsewhere that *Balaustium* mites (Erythraeidae) have very generalized feeding habits, including references to attacks on humans causing dermatitis (Newell 1963, Ido *et al.* 2003).

It appears that all taxa of lice (Anoplura) that affect humans occur in the Cape Verde archipelago, i.e. *Pediculus humanus* (Pediculidae) and *Phthirus pubis* (Phthiridae) (van Harten 1993, Arechavaleta *et al.* 2005). Louse-borne diseases affect all levels of society, but they are most common under poor hygienic circumstances and extreme poverty. Of the two, only *P. pubis* is associated with sexual activity (Brouqui 2011).

Among the 2,000 species of fleas (Siphonaptera) that have been described (Krasnov 2008), at least four occur in the Cape Verde archipelago, i.e. *Ctenocephalides felis*, *Pulex irritans* and *Echidnophaga gallinacea* (Pulicidae) and *Tunga penetrans* (Tungidae) (Gomes 1969, Arechavaleta *et al.* 2005). *P. irritans* and *T. penetrans* have a preference for human blood (Lehane 2005, Krasnov 2008). They are potential vectors of numerous infectious agents, among them viruses and bacteria, and especially *Yersinia pestis*, the causal agent of Black Death (Lehane 2005, Krasnov 2008).

## CONCLUSIONS

Although many bloodsucking arthropods, including known vectors, occur in the Cape Verde Islands, only few studies have been carried out on their biology and role in disease transmission in the archipelago. Most studies have largely or exclusively been carried out by

foreign researchers, illustrating the need to encourage local research teams to study the biology of these species, which include several taxa imposing serious threats to public health, and obtain a better understanding of their environmental requirements in Cape Verde.

## ACKNOWLEDGEMENTS

I would like to express my thanks to Dr Joana Alves (Ministry of Health, Praia, Cape Verde) for providing valuable information and for her

encouragement while I was writing this review. Helpful comments on the manuscript were received from two anonymous referees.

## REFERENCES

- Alves, J., 2004. Contribuição para o estudo de focos residuais de malária em Santiago - Cabo Verde. Master's thesis, Instituto de Higiene e Medicina Tropical, Lisbon.
- Alves, J., B. Gomes, R. Rodrigues, J. Silva, A.P. Arez, J. Pinto & C.A. Sousa, 2010. Mosquito fauna on the Cape Verde Islands (West Africa): an update on species distribution and a new finding. *Journal of Vector Ecology* 35: 307-312.
- Alves, J., A. de Pina, M. Diallo & I. Dia, in press. First report of *Culex tritaeniorhynchus* Giles, 1901 (Diptera, Culicidae) in the Cape Verde Islands. *Zoologia Caboverdiana*.
- Arechavaleta, M., N. Zurita, M.C. Marrero, J.L. Martin (eds.), 2005. Lista preliminar de especies silvestres de Cabo Verde (hongos, plantas y animales terrestres). Conserjería de Medio Ambiente y Ordenación Territorial, Canarias. 155 pp.
- Azevedo, J.F. & H. Moreira, 1946. Um caso de miase interna devida à *Stomoxys calcitrans*. *Anais do Instituto de Medicina Tropical* 3: 467-473.
- Basto, A.P., R.S. Portugal, R.J. Nix, C. Cartaxeiro, F. Boinas, L.K. Dixon, A. Leitão & C. Martins, 2006. Development of a nested PCR and its internal control for the detection of African swine fever virus (ASFV) in *Ornithodoros erraticus*. *Archives of Virology* 151: 819-826.
- Basu, A.K., M. Basu & A. Adesiyun, 2012. A review on ticks (Acari: Ixodoidea: Ixodidae, Argasidae), associated pathogens and diseases of Trinidad and Tobago. *Acarologia* 52: 39-50.
- Benzerroug, B.A., 2005. Mission d'évaluation de la prévalence de la filariose lymphatique. Mission Cap Vert, Novembre 2004-Avril 2005. Organisation Mondiale de la Santé, Programme Mondial d'Élimination de la Filariose Lymphatique. Unpublished report.
- Boorman, J. & A. van Harten, 1992. Vectors of African horse sickness in the Cape Verde Islands. *Veterinary Record* 131: 56.
- Brouqui, P., 2011. Arthropod-borne diseases associated with political and social disorder. *Annual Review of Entomology* 56: 357-374.
- Cambournac, F.J.C., V. Petrarca & M. Coluzzi, 1982. *Anopheles arabiensis* in the Cape Verde archipelago. *Parassitologia* 24: 265-267.
- Cambournac, F.J.C., M.C. Oliveira, A. Correia, M.A. Coutinho, J. Torinho & A.B. Soares, 1984. *Culex (Lutzia) tigripes* (Grandpré); mais uma espécie nova para Cabo Verde. *Anais do Instituto de Higiene e Medicina Tropical* 10: 41-46.
- Chown, S.L. & S.W. Nicolson, 2004. *Insect physiological ecology. Mechanisms and patterns..* Oxford University Press, New York. 256 pp.
- Consoli, R.A. & R. Lourenço de Oliveira, 1994. Principais mosquitos de importância sanitária no Brasil. Editora Fiocruz, Rio de Janeiro. 228 pp.
- Delaunay, P., V. Blanc, P. Del Giudice, A. Levy-Bencheton, O. Chosidow, P. Marty & P. Brouqui, 2011. Bedbugs and infectious diseases. *Clinical Infectious Diseases* 52: 200-210.
- Denion E, P.H. Dalens, P. Couppié, C. Aznar, D. Sainte-Marie, B. Carme, J. Petitbon, R. Pradinaud & M. Gérard, 2004. External ophthalmomyiasis caused by *Dermatobia hominis*. A retrospective study of nine cases and a review of the literature. *Acta Ophthalmologica Scandinavica* 82: 576-584.
- Desquesnes, M. & M.L. Dia 2003a. *Trypanosoma vivax*: mechanical transmission in cattle by one of the most common African tabanids, *Atylotus agrestis*. *Experimental Parasitology* 103: 35-43.
- Desquesnes, M. & M.L. Dia, 2003b. Mechanical transmission of *Trypanosoma congolense* in cattle by the African tabanid *Atylotus agrestis*. *Experimental Parasitology* 105: 226-231.
- Dia, I., J. Alves, A. de Pina, C. Gomes & J.M. Rodriguez, 2011. Mission d'appui sur l'étude de la bio-écologie et la sensibilité aux insecticides des vecteurs du paludisme au Cap-Vert. Unpublished report, Ministry of Health, Praia.
- Dia, I., C.T. Diagne, Y. Ba, D. Diallo, L. Konate & M. Diallo, 2012. Insecticide susceptibility of *Aedes aegypti* populations from Senegal and Cape Verde Archipelago. *Parasites & Vectors* 5:238 (4 pp.); doi:10.1186/1756-3305-5-238.
- Diallo, M., 2003. Mission de consultation entomologique sur les vecteurs du paludisme dans l'île de Santiago (République du Cap-



- Vert). Unpublished report, Ministry of Health, Praia.
- Duarte, E.H., E.E. Correia, C.E. Varela & A. Varela, 2012. Reproduction of mosquitoes (Diptera: Culicidae) in Santa Cruz, Santiago island, Cape Verde Islands. *Zoologia Caboverdiana* 3: 29-36.
- Duarte, E.H., J. Pereira, H. Oliveira, H.S. Lima, A. Perez & E. Pile, 2013. *Aedes (Stegomyia) aegypti* (Diptera: Culicidae) em algumas ilhas de Cabo Verde: Tipologia dos criadouros e sua relação com a presença larval. *Arquivos do Instituto Biológico* 80: 359-362.
- Estrada-Peña, A., A.J. Mangold, S. Nava, J.M. Venzal, M. Labruna & A.A. Guglielmone, 2010. A review of the systematics of the tick family Argasidae (Ixodida). *Acarologia* 50: 317-333.
- FAO, 2000. Recognizing African swine fever. A field manual. FAO Animal Health Manual No. 9. FAO, Rome. 44 pp.
- Franco, A. & A. Menezes, 1955. A filariase autóctone (*W. bancrofti*) na ilha de Santiago (Estudo preliminar). *Anais do Instituto de Medicina Tropical* 12: 369-393.
- Gomes, A., 1969. Algumas espécies de pulgas da ilha de Santiago (Cabo Verde). *Garcia de Orta* 17: 271-274.
- Gomes, B., J. Alves, C.A. Sousa, M. Santa-Ana, I. Vieira, L.T. Silva, A.P. Almeida, M.J. Donnelly & J. Pinto, 2012. Hybridization and population structure of the *Culex pipiens* complex in the islands of Macaronesia. *Ecology and Evolution* 2: 1889-1902.
- Gómez-Díaz, E., J.A. Morris-Pocock, J. González-Solis & K.D. McCoy, 2012. Trans-oceanic host dispersal explains high seabird tick diversity on Cape Verde islands. *Biology Letters* 8: 616-619.
- Götsch, S., M. Leschnik, G. Duscher, J.P. Burgstaller, W. Wille-Piazza & A. Joachim, 2009. Ticks and haemoparasites of dogs from Praia, Cape Verde. *Veterinary Parasitology* 166: 171-174.
- Gratz, N.G., 1999. Emerging and vector-borne resurging diseases. *Annual Review of Entomology* 44: 51-75.
- Grimaldi, D. & M.S. Engel, 2005. *Evolution of the insects*. Cambridge University Press. 772 pp.
- Haitlinger, R., 2009. Three new species of mites (Acari: Prostigmata: Erythraeidae) from the Republic of Cape Verde. *Biologia* 64: 1150-1156.
- Harbach, R.E., 2007. The Culicidae (Diptera): a review of taxonomy, classification and phylogeny. *Zootaxa* 1668: 591-638.
- Ido, T., M. Kumakiri, L.M. Lao, Y. Yano & N. Takada, 2003. Dermatitis caused by *Balaustium murorum*, *Acta Dermatovenereologica* 84: 80-81.
- Karami, A., 2012. Molecular Biology of *Borrelia burgdorferi*. Pp. 1-26 in: A. Karami, (ed.), Lyme disease. InTech, Rijeka & Shanghai. Available at: <http://www.intechopen.com/books/lyme-disease/-molecular-biology-of-lyme-disease-agent>
- Kettle, D.S., 1962. The bionomics and control of *Culicoides* and *Leptoconops* (Diptera, Ceratopogonidae = Heleidae). *Annual Review of Entomology* 7: 401-418.
- Kettle, D.S., 1977. Biology and bionomics of blood-sucking ceratopogonids. *Annual Review of Entomology* 22: 33-51.
- Kirchner, M., A. Brunner, R. Edelhofer & A. Joachim, 2008. Vector-borne parasites of dogs on the Islands of Cabo Verde. *Wiener Klinische Wochenschrift* 120: 49-53.
- Krasnov, B.R., 2008. Functional and evolutionary ecology of fleas. A model for ecological parasitology. Cambridge University Press. 608 pp.
- Lehane, M.J., 2005. The biology of blood-sucking in insects. 2<sup>nd</sup> edition. Cambridge University Press. 336 pp.
- MacLachlan, N.J. & Guthrie, A.J., 2010. Re-emergence of bluetongue, African horse sickness and other *Orbivirus* diseases. *Veterinary Research* 41:35 (12 pp). Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2826768/pdf/vetres-41-35.pdf>
- Mahunka, S., 1991. New and interesting mites from the Geneva Museum LXX. Oribatids from the Cape Verde Islands II (Acari: Oribatida). *Revue Suisse de Zoologie* 98: 567-580.
- Meira, M.T., J. Ruffié & H.T. Sousa, 1957. Algumas carraças das ilhas de Cabo Verde. *Anais do Instituto de Medicina Tropical* 14: 425-427.
- Mellor, P.S., J. Boorman & M. Baylis, 2000. *Culicoides* biting midges: their role as arbovirus vectors. *Annual Review of Entomology* 45: 307-340.
- Middlekauff, W.W. & R.S. Lane, 1980. Adult and immature Tabanidae (Diptera) of

- California. Bulletin of the California Insect Survey 22: 1-99.
- Monteiro, M.L., 2010. Vigilância epidemiológica. In: Actas do Simpósio Internacional de Reflexão sobre a Epidemia de Dengue em Cabo Verde, Praia, 5-7 de Abril de 2010. CD-Rom.
- Newell, I.M., 1963. Feeding habits in the genus *Balaustium* (Acarina, Erythraeidae), with special reference to attacks on man. Journal of Parasitology 49: 498-502.
- Parola, P. & D. Raoult, 2001. Ticks and tickborne bacterial diseases in humans: an emerging infectious threat. Clinical Infectious Diseases 32: 897-928.
- Penrith, M.L., 1998. Control and eradication of an African Swine Fever (ASF) epizootic in Republic of Cape Verde. Food and Agricultural Organization of the United Nations, Pretoria. Available at: <http://www.fao.org/docrep/field/382977.htm>
- Randolph, S.E., 1998. Ticks are not insects: consequences of contrasting vector biology for transmission potential. Parasitology Today 14: 186-192.
- Rahola N., S.M. Goodman & V. Robert, 2011. The Hippoboscidae (Insecta: Diptera) from Madagascar, with new records from the "Parc National de Midongy Befotaka". Parasite 18: 127-140.
- Ribeiro, H., H.C. Ramos, C.A. Pires & R.A. Capela, 1980. Os mosquitos de Cabo Verde (Diptera: Culicidae). Sistemática, distribuição, bioecologia e importância médica. Estudos, Ensaios e Documentos (Junta de Investigações Científicas do Ultramar) 135: 1-141 + 18 maps, 79 figs.
- Rodriguez, J.M., J.O. Guintran, C. Gomes, S. Fall, A. Rietveld, R. Cibulskis, R.D. Newman & R. Kurdova-Mintcheva, 2012. Moving to malaria elimination in Cape Verde. Malaria Journal 11 (Suppl. 1): 09 Available at: <http://www.malariajournal.com/content/pdf/1475-2875-11-S1-O9.pdf>
- Ruppert, E.E. & R.D. Barnes, 1994. Invertebrate Zoology. 6<sup>th</sup> edition. Saunders College Publishing, Orlando, FL. 1100 pp.
- Sellers, R.F., D.E. Pedgley & M.R. Tucker, 1977. Possible spread of African horse sickness on the wind. Journal of Hygiene 79: 279-298.
- Tendeiro, J., 1954. Nota sobre uma pupípara e um ixodídeo de Cabo Verde - *Hippobosca maculata* Leach 1817 e *Amblyomma variegatum* (Fabricius 1794). Garcia de Orta 2: 199-203.
- Turell, M.J., 2012. Members of the *Culex pipiens* complex as vectors of viruses. Journal of the American Mosquito Control Association 28: 123-126.
- van Harten, A., 1993. Terrestrial arthropods of the Cape Verde Islands. A Check-List. Courier Forschungsinstitut Senckenberg 159: 235-309.
- Vazeille, M., Y. Yébakima, R. Lourenço de Oliveira, B. Andriamahefazafy, A. Correia, J.M. Rodrigues, A. Veiga, A. Moreira, I. Leparç-Goffart, M. Grandadam & A.B. Failloux, 2013. Oral receptivity of *Aedes aegypti* from Cape Verde for yellow fever, dengue, and chikungunya viruses. Vector-Borne and Zoonotic Diseases 13: 37-40.
- Vial, L., 2009. Biological and ecological characteristics of soft ticks (Ixodida: Argasidae) and their impact for predicting tick and associated disease distribution. Parasite 16: 191-202.
- WHO, 2004. Global Strategic Framework for Integrated Vector Management. World Health Organization, Geneva. 12 pp. Available at: [http://whqlibdoc.who.int/hq/2004/WHO\\_CD\\_S\\_CPE\\_PVC\\_2004\\_10.pdf](http://whqlibdoc.who.int/hq/2004/WHO_CD_S_CPE_PVC_2004_10.pdf)
- WHO, 2009. Dengue fever, Cape Verde. Weekly Epidemiological Record 84 (45): 469. Available at: [http://www.who.int/wer/2009/wer\\_8445.pdf](http://www.who.int/wer/2009/wer_8445.pdf)
- WHO, 2012. Eliminating malaria. Moving towards sustainable elimination in Cape Verde. World Health Organization, Geneva. 45 pp. Available at: <http://globalhealthsciences.ucsf.edu/sites/default/files/content/ghg/mei-eliminating-malaria-cape-verde-lowres.pdf>
- Zimmer, J.Y., B. Losson & E. Haubruge, 2008. Biologie et écologie des culicoïdes, vecteur de la fièvre catarrhale ovine. Entomologie faunistique 61: 53-57.
- Zumpt, F., 1973. The Stomoxyine biting flies of the world. Diptera; Muscidae. Taxonomy, biology, economic importance and control measures. Gustav Fischer Verlag, Stuttgart. viii + 175 pp.

## APPENDIX 1. Haematophagous arthropods of the Cape Verde Islands and sources used in this review.

<b>Taxonomic group</b>	<b>Authors</b>	<b>Source</b>	<b>Taxa</b>	<b>Researchers (Cape Verde and/or Foreign)</b>
Mosquitoes (Insecta: Nematocera: Culicidae)	Ribeiro <i>et al.</i> (1979)	Journal	<i>Anopheles gambiae s.l.</i>	Foreign
	Ribeiro <i>et al.</i> (1980)	Book	<i>Aedes aegypti</i> , <i>Ae. caspius meirais</i> , <i>Anopheles gambiae s.l.</i> , <i>An. pretoriensis</i> , <i>Culex bitaeniorynchus</i> (Syn. <i>Cx. ethiopicus</i> ), <i>Cx. p. pipiens</i> , <i>Cx. p. quinquefasciatus</i> , <i>Culiseta longiareolata</i>	Foreign
	Cambournac <i>et al.</i> (1982)	Journal	<i>Anopheles arabiensis</i>	Foreign
	Cambournac <i>et al.</i> (1984)	Journal	<i>Culex tigripes</i>	Foreign/CV
	Diallo (2003)	Report	<i>Anopheles arabiensis</i> , <i>An. pretoriensis</i> , <i>Aedes aegypti</i> , <i>Culex pipiens s.l.</i>	Foreign
	Alves <i>et al.</i> (2010)	Journal	<i>Aedes aegypti</i> , <i>Ae. caspius</i> , <i>Anopheles arabiensis</i> , <i>An. pretoriensis</i> , <i>Culex bitaeniorynchus</i> (Syn. <i>Cx. ethiopicus</i> ), <i>Cx. perexiguus</i> , <i>Cx. p. pipiens</i> , <i>Cx. p. quinquefasciatus</i> , <i>Cx. pipiens s.l.</i> hybrids, <i>Cx. tigripes</i> , <i>Culiseta longiareolata</i>	CV/Foreign
	Dia <i>et al.</i> (2011)	Report	<i>Anopheles arabiensis</i> , <i>An. pretoriensis</i> , <i>Aedes aegypti</i> , <i>Ae. caspius</i> , <i>Culex bitaeniorynchus</i> (Syn. <i>Cx. ethiopicus</i> ), <i>Cx. pipiens s.l.</i> , <i>Culex sp.</i>	Foreign/CV
	Dia <i>et al.</i> (2012)	Journal	<i>Aedes aegypti</i>	Foreign
	Duarte <i>et al.</i> (2012)	Journal	<i>Aedes aegypti</i> , <i>Anopheles gambiae s.l.</i> , <i>Culex sp.</i>	CV
	Vazeille <i>et al.</i> (2012)	Journal	<i>Aedes aegypti</i> ssp. <i>formosus</i>	Foreign/CV
	Duarte <i>et al.</i> (2013)	Journal	<i>Aedes aegypti</i>	CV/Foreign
	Alves <i>et al.</i> in press	Journal	<i>Culex tritaeniorhynchus</i>	CV/Foreign

## APPENDIX 1 continued.

	Tendeiro <i>et al.</i> (1954)	Journal	<i>Amblyomma variegatum</i>	Foreign
Ticks (Arachnida: Ixodida)	Meira <i>et al.</i> (1957)	Journal	<i>Amblyomma variegatum</i> , <i>Hylomma sp.</i> , <i>Margaropus decoloratus</i> , <i>Rhipicephalus sanguineos</i>	Foreign
	Kirchner <i>et al.</i> (2008)	Journal	<i>Rhipicephalus sanguineos</i>	Foreign
	Götsch <i>et al.</i> (2009)	Journal	<i>Rhipicephalus sanguineos</i>	Foreign
	Gómez-Díaz <i>et al.</i> (2012)	Journal	<i>Ornithodoros capensis</i>	Foreign
Cattle flies (Insecta: Brachycera: Muscidae)	Arechavaleta <i>et al.</i> (2005)*	Book	<i>Stomoxys calcitrans</i> , <i>S. sitiens</i> , <i>S. niger</i>	Foreign
Horse flies (Insecta: Brachycera: Tabanidae)	Arechavaleta <i>et al.</i> (2005)	Book	<i>Atylotus agrestis</i>	Foreign
Biting midges (Insecta: Nematocera: Ceratopogonida)	Boorman & van Harten (1992)	Journal	<i>Culicoides imicola</i> , <i>C. schultzei</i> , <i>C. nivosus</i>	Foreign
Other taxa	van Harten (1993)	Journal	<i>Simulium rificorne</i> , <i>Ctenocephalides felis</i> , <i>Echidnophaga galinacea</i> , <i>Pulex irritans</i> , <i>Tunga penetrans</i> , <i>Pediculus humanus</i> , <i>Pthirus pubis</i> , <i>Cimex hemipterus</i> , <i>Hippobosca rufipes</i> , <i>H. equina</i> , <i>Olfersia aenescens</i> , <i>Oestrus ovis</i>	Foreign
	Haitlinger (2009)	Journal	<i>Leptus salicus</i> , <i>L. korneli</i> , <i>Erythraeus capeverdensis</i>	Foreign

Received 14 August 2013  
Revision received 4 March 2014  
Accepted 25 April 2014